

**BEST AVAILABLE COPY**

**PCT/EP200 4 / 0 0 6 8 0 5**



**Europäisches  
Patentamt**

**European  
Patent Office**

**Office européen  
des brevets**

**REC'D 20 AUG 2004**

**WIPO**

**PCT**

**Bescheinigung**

**Certificate**

**Attestation**

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

**Patentanmeldung Nr. Patent application No. Demande de brevet n°**

**03394056.0**

**PRIORITY  
DOCUMENT**

**SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)**

**Der Präsident des Europäischen Patentamts;  
Im Auftrag**

**For the President of the European Patent Office**

**Le Président de l'Office européen des brevets  
p.o.**

**R C van Dijk**

Anmeldung Nr:  
Application no.: 03394056.0  
Demande no:

Anmeldetag:  
Date of filing: 20.06.03  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

NESTEC S.A.  
Avenue Nestlé 55  
CH-1800 Vevey  
SUISSE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se referer à la description.)

Modulation of coffee flavour precursor levels in green coffee grains

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

G12N9/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PT RO SE SI SK TR LI

## Modulation of Coffee Flavour Precursor Levels in Green Coffee Grains

### Background of the Invention:

Coffee contains a highly complex mixture of flavour molecules. Extensive research on the composition of instant and fresh ground coffee beverages has, to date, identified more than 850 compounds, many of which are flavour active molecules (Flament, I (2002) Coffee Flavor Chemistry, John Wiley and Sons, UK). However, few of the final coffee flavour molecules found in the cup are present in the raw material, the green grain (green beans) of the plant species *Coffea arabica* or *Coffea canephora (robusta)*. In fact, the majority of the coffee flavour compounds are generated during one or more of the multiple processing steps that occur from the harvest of the ripe red coffee cherries to the final roasted ground coffee product, or extracts thereof, for example soluble coffee products.

The various steps in the production of coffee are described in Smith, A.W., in Coffee; Volume 1: Chemistry pp 1-41, Clark, R.J. and Macrea, R. eds, Elsevier Applied Science London and New York, 1985; Clarke, R.J., in Coffee: Botany, Biochemistry, and Production of Beans and Beverage, pp 230-250 and pp 375-393; and Clifford, M.N. and Willson, K.C. eds, Croom Helm Ltd, London. Briefly, the process starts with the collection of mature, ripe red cherries. The outer layer, or pericarp, can then be removed using either the dry or wet process. The dry process is the simplest and involves 1) classification and washing of the cherries, 2) drying the cherries after grading (either air drying or mechanical drying), and 3) dehusking the dried cherries to remove the dried pericarp. The wet process is slightly more complicated, and generally leads to the production of higher quality green beans. The wet process is more often associated with *C. arabica* cherries. The wet process consists of 1) classification of the cherries, 2) pulping of the cherries, this step is done soon after harvest and generally involves mechanical removal of the "pulp", or pericarp, of the mature cherries, 3) "fermentation", the mucilage that remains attached to the grain of the cherries after pulping is removed by allowing the grain plus attached mucilage to be incubated with water in tanks using a batch process. The "fermentation" process is allowed to continue up to 80 hours, although often 24 hours is generally enough to allow an acceptable fermentation and to cause the pH to drop from around 6.8-6.9 to 4.2-4.6, due to various enzymatic activities and the metabolic action of microorganisms which grow during the fermentation, 4) drying, this

step involves either air or mechanical hot air drying of the fermented coffee grain and 5) “hulling”, this step involves the mechanical removal of the “parch” of the dried coffee grain (dried parchment coffee) and often the silverskin is also removed at this stage. After wet or dry processing, the resulting green coffee grain are often sorted, with most sorting procedures  
5 being based on grain size and/or shape.

The next step in coffee processing is the roasting of the green grain after dehusking or dehulling of dry or wet processed coffee, respectively. This is a time-dependent process which induces significant chemical changes in the bean. The first phase of roasting occurs  
10 when the supplied heat drives out the remaining water in the grain. When the bulk of the water is gone, roasting proper starts as the temperature rises towards 190-200°C. The degree of roasting, which is usually monitored by the colour development of the beans, plays a major role in determining the flavour characteristics of the final beverage product. Thus, the time and temperature of the roasting are tightly controlled in order to achieve the desired coffee  
15 flavour profile. After roasting, the coffee is ground to facilitate extraction during the production of the coffee beverage or coffee extracts (the latter to be used to produce instant coffee products). Again, the type of grinding can influence the final flavour of the beverage.

While a considerable amount of research has been carried out on the identification of the  
20 flavour molecules in coffee, much less work has been done regarding the physical and chemical reactions which occur within the coffee grains during each of the processing steps. This latter point is particularly evident for the roasting reaction, where the large number of grain constituents undergo an extremely complex series of heat induced reactions (Homma, S. 2001, In “Coffee: Recent Developments”. R.J. Clarke and O.G. Vitzthum eds, Blackwell  
25 Science, London; Yeretdzian, C., *et al* ((2002) Eur. Food Res. Technol. 214, 92-104; Flament, I (2002) Coffee Flavor Chemistry, John Wiley and Sons, UK; Reineccius, G.A., “The Maillard Reaction and Coffee Flavor” Conference Proceedings of ASIC, 16<sup>th</sup> Colloque, Kyoto, Japan 1995). Because of the large number of the potential reactants, and the complexity of the reactions involved, little research data currently exist which establishes  
30 strong links between specific biological molecules present in the green beans and the numerous flavour molecules identified in a cup of coffee.

Nonetheless, while the details of most of the reactions that occur during the different steps of coffee processing remain relatively unclear, it is thought that an important flavour generating

reaction responsible for many of the flavours associated with coffee aroma is the "Maillard" reaction during coffee roasting. A vigorous Maillard reaction occurs between the grain reducing sugars/polysaccharide degradation products and the amino group containing molecules (particularly the proteins, peptides, and amino acids) during the roasting step.

Because the Maillard reaction apparently makes an important contribution to the generation of coffee flavour and aroma molecules during coffee roasting, there might be an association between the levels of primary Maillard reactants in the green beans and the quality of the flavour/aroma developed after roasting.

As noted above, an important group of substrates in the Maillard reaction are the amino acids, peptides and proteins. To date, there are no firm associations between the levels, or types, of amino group containing molecules in different coffees and the qualities of those coffees.

Although the amino acid contents of *arabica* and *robusta* coffee beans have been analysed (M.N. Clifford 1985, In "Coffee: Botany, Biochemistry, and Production of Beans and Beverage", Clifford and Willson eds, Croom Helm London; Arnold *et al* 1996, Z. Lebensm Unters Forsch., Vol 199, 22-25; Ludwig *et al* 2000, Eur. Food Res Technol., Vol 211, 111-116.), there are no reports directly linking specific levels or ratios of amino acids and high or low flavour qualities. Using 2-D electrophoresis, it has been shown that differences exist in the levels and amounts of the major storage proteins in *arabica* and *robusta* green coffee beans – however, no association between these storage protein differences and flavour quality was noted (Rogers *et al*, 1999, Plant Physiol. Biochem. Vol 37, 261-272). It has also recently been found that small differences exist between the storage proteins of immature and mature coffee beans, which have different flavour qualities (Montavon, P. *et al*, 2003, J. Agric and Food Chemistry Vol 51, 2328-2334). However, because there are many changes occurring during seed maturation, this latter work only suggests a link may exist between the quality improvement caused by maturation and the differences seen in the 2-D gel patterns of the main coffee storage proteins. Overall, currently no clear evidence exists linking any differences seen for the coffee storage proteins, or other major green bean proteins, and the flavour qualities of coffee.

It has recently been shown that there are differences in the profiles of peptides isolated from *arabica* and *robusta* green beans (Ludwig *et al* 2000, Eur. Food Res Technol., Vol 211, 111-116.). Although their results showed that the *arabica* and *robusta* peptide extracts differ in

their aroma precursor profile, the data presented in this report do not identify which component(s) in the extracts is / are responsible for these aroma profile differences. These workers also detected at least two different proteinase activities in crude extracts of the green coffee, but they did not correlate any specific activities with aroma/flavour quality (Ludwig *et al* 2000, Eur. Food Res Technol., Vol 211, 111-116). Finally, it is also thought that the very high temperatures used during the later stages of green coffee grain roasting cause substantial cleavage of the proteins present in the coffee grain (Homma, S. 2001, In "Coffee: Recent Developments". R.J. Clarke and O.G. Vitzthum eds, Blackwell Science, London; Montavon, P., *et al* 2003, "Changes in green coffee protein profiles during roasting", J. Agric. Food Chem. 51, 2335-2343). However, the overall scheme for this protein degradation is very poorly understood, but presumably depends on, among other things, the precise state of the main coffee proteins in the raw material before the start of roasting. To our knowledge, there are no other significant reports addressing the possibility that peptide profiles in coffee could be involved in the production of coffee aroma/flavour.

In the roasting of the fermented seeds of *Theobroma cacao* (cocoa beans), there would appear to be an involvement of seed amino acids and peptides in the development of Maillard reaction aromas/flavours. Relative to other seeds, *T. cacao* seeds have been shown to have an unusually high level of aspartic proteinase activity (Biehl, B., Voigt, J., Voigt, G., Heinrichs, H., Senyuk, V. and Bytof, G. (1994) "pH dependent enzymatic formation of oligopeptides and amino acids, the aroma precursors in raw cocoa beans". In *The Proceedings of the 11<sup>th</sup> International Cocoa Research Conference*, 18-24 July 1993, Yamoussoukro, Ivory Coast). In order to produce cocoa beans with a high level of cocoa flavour precursors, it is necessary to carry out a natural fermentation step (unfermented beans develop little flavour when roasted). During this fermentation step, the sugars in the pulp are fermented, generating high levels of acids, particularly acetic acid (Carr, J.G. (1982) Cocoa. In *Fermented Foods*. Economic Microbiology. Vol 7. pages 275-292. (A.H. Rose ed). Academic Press). As the fermentation continues, the pH in the seed decreases and the cell structure becomes disrupted. The low pH triggers the abundant cacao seed aspartic proteinase to become mobilized and/or activated, resulting in a massive degradation of cellular protein (Biehl, B., Passern, D., and Sagemann, W. (1982) "Effect of Acetic Acid on Subcellular Structures of Cocoa Bean Cotyledons". J. Sci. Food Agric. 33, 1101-1109; Biehl, B., Brunner, E., Passern, D., Quesnel, V.C., and Adomako, D. (1985) "Acidification, proteolysis and flavour potential in fermenting cocoa beans". J. Sci. Food Agric. 36, 583-598). Peptides and amino acids have

been shown to be cocoa flavour precursors ( Rohan, T. (1964) "The precursors of chocolate aroma: a comparative study of fermented and unfermented cocoa beans". J. Food Sci., 29, 456-459; Voigt, J. and Biehl, B. (1995) "Precursors of the cocoa specific aroma components are derived from the vicilin-class (7S) globulin of the cocoa seeds by proteolytic processing". Bot. Acta 108, 283-289). Thus, the *T. cacao* seed aspartic proteinase, together with a seed serine carboxypeptidase, have been proposed to be critical for the generation of cocoa flavour precursors during fermentation (Voigt, J. and Biehl, B. (1995) "Precursors of the cocoa specific aroma components are derived from the vicilin-class (7S) globulin of the cocoa seeds by proteolytic processing". Bot. Acta 108, 283-289; Voigt, J., Heinrichs, H., Voigt, G. and Biehl, B. (1994) "Cocoa-specific aroma precursors are generated by proteolytic digestion of the vicilin-like globulin of cocoa seeds". Food Chemistry, 50, 177-184.) The gene encoding the abundant cacao seed aspartic proteinase has been identified and a method to over-express this protein in cacao seeds which can generate increased levels of cacao flavour precursor amino acids and peptides in fermented cocoa beans has recently been described in International Patent Publication No. 02/04617, the whole contents of which are incorporated herein by reference. However, the teaching of International Patent Publication No. 02/04617 is directed towards cacao seeds, which undergo a specific long acid fermentation step, unlike coffee grains which do not.

## Objects of the Invention

It is an object of the present invention to improve the flavour quality of coffee.

More specifically, it is an object of the present invention to improve the levels of the flavour precursors in the raw material (the green grain) so that, following post harvest treatment and roast-processing, an improved flavour is achieved. Without being bound by theory, it is believed that, if there are variations in the levels of peptides and protein degradation between coffees with significantly different flavours, then it is possible that these variations could be due to differences in the endogenous proteinase activities in these different grains. This difference might be detectable at the level of mRNA expression by variations in the levels of expression for particular seed proteinase genes.

## Statements of the Invention

The present invention involves, therefore, identifying gene sequences encoding for coffee grain (seed) specific proteinases and showing that there are indeed variations in the expression of these genes in *arabica* and *robusta*.

- 5 More specifically, the present invention discloses a major coffee cysteine proteinase (CcCP-1), a major coffee cysteine proteinase inhibitor (CcCPI-1) and coffee aspartic proteinases (CcAP-1 and CcAP-2), all of which are expressed in coffee seeds. We further show how either over-expression of these proteins specifically late in seed development, or the reduced expression of these proteins during late seed development, can alter the amino
- 10 acid/peptide/protein profile of the mature beans. By using one or more of the disclosed gene sequences and gene constructs to alter the amino acid/peptide/protein profile of the mature beans, we disclose a new method to alter the flavour precursor profile of mature coffee beans, and thus thereby allow the production of roasted coffee beans with altered flavours.
- 15 In a first aspect, the present invention provides an isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide having cysteine proteinase activity, wherein the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID No. 2 have at least 70%, preferably at least 80%, sequence identity based on the ClustalW alignment method; or the complement of the nucleotide sequence, wherein the complement contains the
- 20 same number of nucleotides as the nucleotide sequence, and the complement and the nucleotide sequence are 100% complementary. Preferably, the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID No. 2 have at least 85%, preferably at least 90%, optionally at least 95%, sequence identity based on the ClustalW alignment method. Preferably, the nucleotide sequence comprises the nucleotide sequence of SEQ ID
- 25 No. 1. Preferably, the polypeptide comprises the amino acid sequence of SEQ ID No. 2.

- In a second aspect, there is provided an isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide having cysteine proteinase inhibitor activity, wherein the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID No. 4 have
- 30 at least 70%, preferably at least 80%, sequence identity based on the ClustalW alignment method; or the complement of the nucleotide sequence, wherein the complement contains the same number of nucleotides as the nucleotide sequence, and the complement and the nucleotide sequence are 100% complementary. Preferably, the amino acid sequence of the polypeptide and the amino acid sequence of SEQ ID No. 4 have at least 85%, preferably at



least 90%, optionally at least 95%, sequence identity based on the ClustalW alignment method. Preferably, the nucleotide sequence comprises the nucleotide sequence of SEQ ID No. 3. Preferably, the polypeptide comprises the amino acid sequence of SEQ ID No. 4.

- 5 In a third aspect, there is provided an isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide having aspartic endoproteinase activity, wherein the amino acid sequence of the polypeptide and the amino acid sequence selected from SEQ ID No. 6 or 8, preferably SEQ ID No. 8, have at least 75%, preferably at least 80%, sequence identity based on the ClustalW alignment method, or the complement of the nucleotide sequence,
- 10 wherein the complement contains the same number of nucleotides as the nucleotide sequence, and the complement and the nucleotide sequence are 100% complementary. Preferably, the amino acid sequence of the polypeptide and the amino acid sequence selected from SEQ ID No. 6 or 8, preferably SEQ ID No. 8, have at least 85%, preferably at least 90%, optionally at least 95%, sequence identity based on the ClustalW alignment method. Preferably, the
- 15 nucleotide sequence comprises the nucleotide sequence of SEQ ID No. 5 or 7, preferably SEQ ID No. 7. Preferably, the polypeptide comprises the amino acid sequence of SEQ ID No. 6 or 8, preferably SEQ ID No. 8.

- 20 In a further aspect, there is provided a vector comprising the polynucleotide of any one of first to third aspects of the invention.

- 25 In a further aspect, there is provided a recombinant DNA construct comprising the polynucleotide of any one of first to third aspects of the invention, operably linked to a regulatory sequence.

- In a further aspect, there is provided a method for transforming a cell comprising transforming the cell with the polynucleotide of any one of first to third aspects of the present invention.

- 30 In a further aspect, there is provided a cell comprising the aforementioned recombinant DNA construct, which cell is preferably a prokaryotic cell, an eukaryotic cell or a plant cell, preferably a coffee cell.

In a further aspect, there is provided a transgenic plant comprising such a transformed cell.

In the present application, coffee cherry terms are defined as follows: coffee cherry; entire fruit; exocarp, skin; pericarp, fleshy major outer layer of cherry; and grain, coffee seed. For a fuller explanation of these terms, reference is made to Clarke, R.J., in Coffee: Botany, Biochemistry, and Production of Beans and Beverage, pp 230, Clifford, M.N. and Willson, K.C. eds, Croom Helm Ltd, London, the contents of which are incorporated in their entirety.

#### Brief Description of the Invention

The invention can be understood from the following detailed description and the accompanying Sequence Listing which forms part of the present application.

Table 1 hereunder lists the polypeptides that are described herein, along with the corresponding sequence identifier (SEQ ID No) as used in the attached listing.

#### **Table 1:**

SEQ ID No 1 (CcCP1 : Cysteine proteinase, nucleic acid and its corresponding amino acid)

SEQ ID No 2 (CcCP1: Cysteine proteinase, amino acid)

SEQ ID No 3 (CcCPI-1 : Cysteine proteinase Inhibitor, nucleic acid and its corresponding amino acid)

SEQ ID No 4 (CcCPI-1: Cysteine proteinase Inhibitor, amino acid)

SEQ ID No 5 (CcAP1 : Aspartic endoproteinase 1, nucleic acid and its corresponding amino acid)

SEQ ID No 6 (CcAP1 : Aspartic endoproteinase 1, amino acid)

SEQ ID No 7 (CcAP2 : Aspartic proteinase 2, nucleic acid and its corresponding amino acid)

SEQ ID No 8 (CcAP2 : Aspartic proteinase 2, amino acid)

The sequence listing employs the one letter codes for nucleotide sequence characters and the three letter codes for amino acids as defined for IUPAC-IUBMB Standards and as described in Nucleic Acids Research 13:3021-3030 (1985), which is incorporated herein by reference.

## Drawings

In the drawings,

5 **Figure 1** shows a Northern blot analysis of cysteine proteinase gene in different tissues of *Coffea arabica*, in which the lanes are labeled R: root, S: stem, L: young leaves; and SG, LG, Y and Red are grain from small green fruit, large green fruit, yellow fruit and red fruit, respectively. Five micrograms of total RNA was loaded in each lane. MW is an RNA size ladder. Panel A illustrates an autoradiography after 24 hours exposure and Panel B  
10 demonstrates the ethidium bromide staining of the gels prior to blotting;

**Figure 2** shows a Northern blot analysis of Cysteine proteinase inhibitor gene in different tissues of *Coffea arabica*, in which the lanes are labeled R: root, S: stem, L: young leaves and SG, LG, Y and Red for grain from small green fruit, large green fruit, yellow fruit and red  
15 fruit, respectively. Five micrograms of total RNA was loaded in each lane. MW is an RNA size ladder. Panel A illustrates an autoradiography after 24 hours exposure and panel B demonstrates the ethidium bromide staining of the gels prior to blotting;

**Figure 3** shows a Northern blot analysis of Cysteine proteinase inhibitor gene in different  
20 stages of development of *Coffea arabica* (ARA) and *Coffea robusta* (ROB) fruit. The lanes are labeled small green fruit (SG), large green fruit (LG), yellow fruit (Y) and red fruit (Red), respectively. Five micrograms of total RNA was loaded in each lane. MW is an RNA size ladder. Panel A illustrates an autoradiography after 24 hours exposure. Panel B demonstrates the ethidium bromide staining of the gels prior to blotting; and  
25

Figure 4 shows a Northern blot analysis of aspartic proteinase 2 (CcAP2) gene in different tissues of *Coffea arabica*, in which the lanes are labelled R: root, S: stem, L: young leaves, F: flowers; SG(G) and (P), LG(G) and (P), Y(G) and (P) and Red(G) and (P) are for grain and for pericarp, respectively, from small green, large green, yellow and red cherries, and SG(G), LG(G), Y(G) and R(G) for pericarp from small green, large green, yellow and red cherries respectively. Five micrograms of total RNA was loaded in each lane. Panel A demonstrates the ethidium bromide staining of large ribosomal RNA prior to blotting as a loading control and panel B is an autoradiogram showing the appearance of the CcAP2 mRNA in the specific tissues tested.

#### Detailed Description of the Invention

As used herein, a “polynucleotide” is a nucleotide sequence such as a nucleic acid fragment. A polynucleotide may be a polymer of RNA or DNA that is single- or double-stranded, that optionally contains synthetic, non-natural or altered nucleotide bases. A polynucleotide in the form of a polymer of DNA may comprise one or more segments of cDNA, genomic DNA, synthetic DNA or mixtures thereof.

Similar nucleic acid fragments are characterised, in the present invention, by the percent identity of the amino acid sequences that they encode, to the amino acid sequences disclosed herein, as determined by algorithms commonly used by those skilled in the art. Suitable nucleic acid fragments (or isolated polynucleotides of the first to third aspects of the present invention) encode polypeptides that are at least 70% identical, preferably at least 80% identical, to the amino acid sequences disclosed herein. Preferred nucleic acid fragments encode amino acid sequences that are at least 85% identical to the amino acid sequences disclosed herein. More preferred nucleic acid fragments encode amino acid sequences that are at least 90% identical to the amino acid sequences disclosed herein. Still more preferred are nucleic acid fragments that encode amino acid sequences that are at least 95% identical to the amino acid sequences disclosed herein. Multiple alignment of sequences should be performed using the ClustalW method of alignment (Thompson *et al*, 1994, Nucleic Acids Research, Vol 22, p4673-4680; Higgins & Sharp 1989 Cabios. 5:151-153).

As used herein, the term “similar nucleic acid fragments” refers to polynucleotide sequences in which changes in one or more nucleotide bases result in substitution of one or more amino

acids, but which changes either do not affect the function of the polypeptide encoded by the nucleotide sequence or do not affect the ability of nucleic acid fragment to mediate gene expression by gene silencing via, for example, antisense or co-expression technology. The term "similar nucleic acid fragments" also refers to modified polynucleotide sequences, in which one or more nucleotide bases is / are deleted or inserted, provided that the modifications do either do not affect the function of the polypeptide encoded by the nucleotide sequence or do not affect the ability of nucleic acid fragment to mediate gene expression by gene silencing. It will, therefore, be understood that the scope of the present invention extends beyond the polynucleotide and polypeptide sequences specifically disclosed herein.

Similar nucleic acid fragments may be selected by screening nucleic acid fragments in the form of subfragments or modified nucleic acid fragments, for their ability to affect the level of the polypeptide encoded by the unmodified nucleic acid fragments in the plant or plant cell.

The term "operably linked" refers to the association of two or more nucleic acid fragments on a single nucleic acid fragment so that the function of one is affected by the other.

"Regulatory sequences" refer to nucleotide sequences located upstream, within, or downstream, of a coding sequence and which influence transcription, RNA processing or stability, or translation of the coding sequence associated therewith. Regulatory sequences may include promoters, translation leader sequences, introns and polyadenylation recognition sequences. When a regulatory sequence in the form of a promoter is operably linked to a coding sequence, the regulatory sequence is capable of affecting the expression of the coding sequence. Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term "expression" refers to the transcription, and stable accumulation, of sense RNA (mRNA) or antisense RNA derived from the nucleic acid fragments of the present invention. Expression may also refer to the translation of mRNA into a polypeptide. Overexpression refers to the production of a gene product in a transgenic cell, that exceeds the level of production in normal, or non-transformed, cells. "Altered levels" refers to the production of gene product(s) in a transgenic cell in amounts or proportions that differ from that of normal, or non-transformed, cells.

“Transformation” refers to the transfer of a nucleic acid fragment into the genome of a host cell, resulting in genetically stable inheritance. Host cells containing the transformed nucleic acid fragments are referred to herein as “transgenic cells.”

5

Standard recombinant DNA and molecular cloning techniques as used herein are well known in the art and are described more fully in Sambrook *et al* "Molecular Cloning: A Laboratory Manual"; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989, which is incorporated herein by reference.

10

### Examples

The following Examples illustrate the invention without limiting the invention to the same. In the examples, all parts and percentages are by weight and degrees are in Celsius, unless this is otherwise specified.

15

In the following Examples, these abbreviations have been used:

PCR : Polymerase chain reaction

RACE : Rapid amplification cDNA ends

20

From the above discussion and the Examples below, those skilled in the art can ascertain the essential features of the present invention, and without departing from the scope thereof can make various changes and modifications thereto, to adapt it to various usages and conditions as desired.

25

### Production of cDNA libraries and screening

#### Production of Seed Specific RNA

Coffee cherries of the *Robusta* variety Q121 were harvested 30 WAF (weeks after flowering) at the ICCRI, Indonesia. The pericarps of these cherries were then removed and the remaining perisperm/endosperm material was frozen and ground to a powder in liquid nitrogen. The RNA was extracted from the frozen powder material using the method described previously for the RNA extraction of cacao seeds (Guilloteau, M. *et al*, 2003, Oil bodies in *Theobroma cacao* seeds: cloning and characterisation of cDNA encoding the 15.8

30

and 16.9 kDa oleosins. Plant Science Vol 164, 597-606). Poly A<sup>+</sup> RNA was prepared from approximately 250µg total RNA using the "PolyA Purist™" kit of AMBION (manufactured by Ambion, Inc.) according to their kit instructions.

#### 5 *Production of First Set of Seed cDNA clones*

Approximately 50-100ng of this poly A<sup>+</sup> RNA was then employed in the synthesis of the first strand cDNA using "SuperScript™ II RNase H<sup>-</sup> reverse transcriptase (GIBCOBRL™) and the SMART™ PCR cDNA synthesis kit (Clontech) as follows. A reaction containing 2µl of 30 WAF poly A<sup>+</sup> RNA, 1 µL CDS oligo (SMART™ PCR cDNA kit, Clontech), 1 µL Smart II oligo (SMART™ PCR cDNA kit, Clontech), and 8 µL deionised H<sub>2</sub>O. This mixture was heated to 72°C for 5 minutes and then placed on ice. Then the following was added; 1 µL 10 mM dNTPs, 4 µL SuperScriptII™ 1<sup>st</sup> stand buffer and 2 µL DTT. This mixture was put at 42°C for 2 minutes then 1 µL of SuperScriptII™ RNaseH<sup>-</sup> reverse transcriptase (200 units/µL GIBCO BRL™) was added and the mixture was incubated in an air circulating incubator at 42°C for a further 50 minutes.

After the reverse transcription reaction, the following PCR reaction was carried out. 98 µL of the Master Mix described in the SMART™ PCR cDNA kit (Clontech) containing Advantage™ 2 polymerase (Advantage™ 2 PCR kit, ClonTech) was set up on ice and then 3 µL of the 1<sup>st</sup> strand cDNA synthesis reaction described above was added. This 100 µL PCR reaction was then placed in a MJ Research PTC-150 HB apparatus and the following PCR conditions were run: 95°C for 1 minute, then 16 cycles of 95°C for 15 seconds, 65°C for 30 seconds, 68°C for 6 minutes. The amplified DNA was purified using the Strataprep™ PCR Purification Kit (Stratagene) according to the suppliers' instructions. The DNA, which was eluted in 50 µL deionized water, was then "polished" using the Pfu-1 polymerase reagents contained in the PCR-Script™ Amp cloning kit (Stratagene) as follows; 50 µL DNA, 5 µL 10 mM dNTPs, 6.5 µL 10 x Pfu-1 polishing buffer, 5 µL cloned Pfu-1 DNA polymerase (0.5 U/µl). This reaction was then incubated at 72°C for 30 minutes in a PCR apparatus with a heated cover (Perkin Elmer). Using the protocol described in the pPCR-Script™ Amp kit (Stratagene), the polished (blunted) PCR products were ligated into the Srf-1 digested pPCR-Script™ Amp SK(+) vector in the presence of Srf-1 enzyme and the ligation reaction products were transformed into the XL-10 Gold™ Kan ultracompetent E. coli cells. Selection for transformation with plasmids containing inserts was done using LB-Amp plates

and IPTG and Xgal spread on the surface as described in the pPCR-Script™ Amp kit. White colonies were selected and the clones were named Dav1-1 etc.

#### *Production of Second Set of seed cDNA clones with Size Selected cDNA*

5 Seeds highly express a small number of proteins, such as the seed storage proteins (White *et al*, 2000, Plant Physiology, Vol 124, 1582-1594). When cDNA is prepared from such tissue, the very high level of the storage proteins and other seed specific proteins leads to a high level of cDNA “redundancy”, that is, the population of cDNA produced contains high proportions of the same cDNA. In order to reduce the redundancy of cDNA made from  
10 coffee seed mRNA, and to selectively characterise long and weakly expressed cDNA, a second cDNA cloning strategy was also used. Using the products of the reverse transcriptase reaction described above, the following PCR reactions was set up using the Advantage™ 2 PCR kit (Clontech): 3 µL of the reverse transcriptase reaction, 5 µL 10 x Advantage™ 2 PCR buffer, 1 µL dNTP's (10 mM each), 2 µL PCR primer (SMART™ PCR cDNA kit, Clontech), 39 µL deionised water, and 1 µL 50 x Advantage™ 2 polymerase mix. This PCR  
15 reaction was then placed in a MJ Research PTC-150 HB apparatus and the following PCR conditions were run: 95°C for 1 minute, then 16 cycles of 95°C for 15 seconds, 65°C for 30 seconds, 68°C for 6 minutes. At the end of the PCR, 1 µL 10% SDS was added with gel loading buffer, the sample was heated to 37°C for ten minutes. The sample was then split for  
20 loading onto a 0.7% agarose gel without ethidium bromide: 10% was loaded into a small well beside a DNA marker lane and the other 90% was loaded into a neighbouring large, preparation scale well. After the gel was run, the gel section with the size markers, plus the 10% reaction sample, were stained with ethidium bromide. This stained gel section was then used as a template to generate gel slices containing PCR amplified cDNA of different sizes  
25 from the cDNA present in the remaining unstained (preparation) part of the gel. Six gel slices were generated having the indicated size range of PCR fragments; A1A (0.8-1kb), A1B (1-1.5 kb), A2 (1.5-2.25 kb), A3 (2.25-3.25), A4 3.25-4 kb), and A5 (4-6.5 kb).

The DNA in each gel slice was eluted from the agarose using the QIAEX II kit from Qiagen  
30 following the suppliers instructions (for samples 3A, 4A, and 5A were heated for 10 minutes at 50°C and 1A, 1B, and 2A were heated for 10 minutes at room temperature). The purified double stranded cDNA was then re-amplified further by PCR with a TAQ enzyme mix which makes fragments having a 3' T overhang as follows: 30 µL of the gel isolated double



stranded cDNA, 5  $\mu$ L 10 x TAQ buffer (supplied with TAQ PLUS precision polymerase mix, Stratagene), 1  $\mu$ L 40 mM dNTP's (each 10 mM), 2  $\mu$ L PCR primer (SMART<sup>TM</sup> PCR cDNA kit, Clontech), 0.5  $\mu$ L TAQ PLUS precision polymerase mix (Stratagene) and 11.5  $\mu$ L deionised water. The PCR reaction conditions were as follows: 95°C for 1 minute then 7 cycles 95°C for 15 seconds, 65°C for 1 minute, 72°C for 8 minutes, then 1 cycle at 95°C for 15 seconds, 65°C for 1 minute, 72°C for 10 minutes.

The PCR amplified DNA produced was then ligated into the vector pCR<sup>TM</sup>-TOPO<sup>TM</sup> and cloned into TOP10 *E. coli* cells using the TOPO<sup>TM</sup> TA kit (Invitrogen) as described by the supplier. The clones were named by their order of isolation and their position in the sizing gel (for example, A2-1, A2-2, etc.).

#### *Seed cDNA Screening and Preliminary Identification*

The first set of white colonies obtained in Dav-1 library were screened by first determining the size of each insert by PCR amplifying the insert using the primers T3 and T7 which flank the cloning site used and examining the PCR amplified fragments on a gel.

Each white colony were resuspended in 200  $\mu$ l sterile water and 10-30  $\mu$ l of this was added to 5  $\mu$ l 10X Taq polymerase buffer (Stratagene), 1  $\mu$ l 10 mM dNTP mix, 2.5  $\mu$ l 20  $\mu$ M T3 primer, 2.5  $\mu$ l 20  $\mu$ M T7 primer, 1  $\mu$ l DMSO, 0.5  $\mu$ l Taq polymerase (Stratagene), and H<sub>2</sub>O up to 50  $\mu$ l final volume). The PCR reaction program uses was 94°C for 1 min, then 30 cycles of 94°C for 1 min, 55°C for 1.5 min and 3.5 min at 72°C, and a final cycle of 7 min at 72°C. To reduce redundancy, the PCR inserts of similar size were subjected to digestion by the restriction enzyme Hae III. Those PCR fragments with the same Hae III restriction pattern were not studied further. The plasmids of clones with PCR fragments >500 bp and which had unique Hae III restriction patterns were then purified by using the Qiawall 8 ultra plasmid kit (Qiagen) for 5' end dideoxy sequencing using the appropriate T7 or T3 sequencing primers coded in the flanking vector sequences. Because the inserts were not cloned in a directed fashion, it was first necessary to determine the 5' end of each clone by a ScaI digestion of the purified plasmid DNA (the CDS SMART primer contains a Sca I site allowing the orientation of the insert to be determined). The DNA sequence data obtained was subsequently blasted against the non-redundant database protein in GENEbank to obtain a preliminary annotation of each cDNA clone using the program BLASTX<sup>TM</sup>.

Seed cDNA banks have a high level of redundancy. That is, a small number of seed mRNA have an unusually high level of expression, such as those encoding the seed storage proteins, and therefore their cDNA are very abundant in seed cDNA banks (White *et al*, 2000, Plant Physiology, Vol 124, 1582-1594). Therefore, as soon as the main redundant cDNA's were identified in the first round of sequencing the coffee seed cDNA, a pre-screening step was added for the white insert containing colonies prior to the determination of insert size. Four sequences were very highly expressed and the following specific primers sets were made for each of these redundant sequences,

- 1) 2S protein, contig 8A 5' AGCAACTGCAGCAAGGTGGAG 3' and contig 8B 5' CGATTTGGCACTGCTGTGGTTC 3' (55°C used in PCR, 114 bp fragment),
- 2) 2S protein contig 15A 5' GCCCGTGCTCCTGAACCA 3' and contig 15B 5' GTATGGTTGCGGTGGCTGAA 3' (55°C used in PCR, 256 bp fragment),
- 3) Oleosin 15.5 contig 30A 5' ACCCCGCTTTTCGTTAT 3' and contig 30B TCTGGCTACATCTTGAGTTCT 3' (55°C used in PCR, 261 bp fragment), and
- 4) 11S protein contig 37A 5' GTTTCAGACCGCCATCAG 3' and contig 37B 5' ATATCCATCCTCTTCCAACACC 3' (59°C used in PCR, 261 bp fragment).

The PCR reactions for this prescreen step were run as follows: 10-30 µl of the white colony in sterile H<sub>2</sub>O, 5 µl 10X Taq buffer (Stratagene), 1 µl 10 mM dNTP, 2.5 µl of each primer at 20 µM, 1 µl DMSO, 0.5 µl Taq polymerase (Stratagene 10U/µl) and sterile H<sub>2</sub>O was added to produce a final reaction total volume of 50 µl. The PCR program was 1 min at 94°C, then 30 cycles of 1 min at 94°C, 1.5 min at specific temperature for each primer pair, 2.5 min at 72°C, followed by 7 min at 72°C.

#### Full Length cDNA Insert Sequencing and Sequence Analysis

cDNA clones whose partial sequences showed initial homologies to proteinases and proteinase inhibitors were fully sequenced on both strands using the standard dideoxy primer walking strategy. The sequences are shown under SEQ ID Nos. 1, 3, 5 and 7. The full length sequences obtained were again blasted against the GenBank non redundant protein database using BLASTX to reinforce the preliminary annotation

Sequence identities of sequence pairs were calculated using the ClustalW™ program contained in the MegAlign™ module of the Lasergene™ software package (DNASTAR Inc). The default parameters were chosen as follows: (1- MULTIPLE ALIGNMENT PARAMETERS - Gap penalty 15.00, Gap length penalty 6.66, Delay divergent Seqs (%) 30, DNA transition weight 0.5, Protein Weight Matrix-Gonnet Series, DNA Weight Matrix IUB. 2- PAIRWISE ALIGNMENT PARAMETERS-Slow/Accurate (Gap Penalty 15.00, Gap Length Penalty 6.66), Protein Weight Matrix-Gonnet 250, DNA Weight Matrix-IUB) and the sequences used were either the full length nucleotide sequence of each cDNA or the full ORF (open reading frame) of each cDNA.

**TABLE 2**

Identity values between the nucleic acid and amino acid sequences of CcCP-1, CcCPI-1, and CcAP-2 and related genes found in the non-redundant protein database of GeneBank and those of WO 02/04617.

cDNA Sequences	nucleotide identity (%)	protein identity (%) (ORF)
CcAP1 vrs TcAP1	2.9	13.3
CcAP1 vrs TcAP2	2.4	9.8
CcAP2 vrs TcAP1	55.0	61.5
CcAP2 vrs TcAP2	55.1	61.3
CcCP-1 vs <i>Arabidopsis thaliana</i> putative cysteine proteinase (AY070063)	51.8	64.3
CcCP-1 vs Glycine max cysteine endopeptidase (Z32795)	49.1	61.3
CcCP-1 vs <i>Vicia sativa</i> cysteine proteinase precursor (Z99172)	49.0	60.9
CcAP2 vs <i>Lycopersicon</i> <i>esculentum</i> aspartic proteinase precursor (L46681)	65.9	71.1
CcAP2 vs <i>Ipomoea batatas</i>		

putative aspartic proteinase mRNA (AF259982)	71.7	69.6
CcAP2 vs <i>Nepenthes alata</i> NaAP4 mRNA for aspartic proteinase 4 (AB045894)	58.4	66.5
CcCPI-1 vs <i>Malus x domestica</i> cystatin (AY176584)	38.8	45.5

### Northern-Blot Analysis

5

Freshly harvested roots, young leaves, stem, flowers and fruit at different stages of development (small green fruit (SG), large green fruit (LG), Yellow fruit (Y) and red fruit (R)) were harvested from *Coffea arabica* var. Catura T-2308 grown under greenhouse conditions (25°C, 70 RH) in Tours, France, and from *Coffea canephora* var. BP409 grown either in Equador or ICCRI, Indonesia. The fresh tissues were frozen immediately in liquid nitrogen and total RNA was isolated from each tissue using the extraction procedure described above. A total of 5 µg of RNA was run on a 1.2% (w/v) denaturing RNA gel containing formaldehyde. The total RNA samples from each plant tissue were heated at 65°C for 15 min in presence of 7 µL "RNA Sample Loading Buffer" (without ethidium bromide, Sigma), and then put immediately on ice for 2 minutes before being loaded onto the 1.2% RNA gel. The gels were run at 60 Volts for 5 hours. The gel was then soaked twice in 10× SSC for 20 min. The RNA in the gel was transferred overnight by capillary transfer to a "Positive TM Membrane" (Qbiogene) in 10× SSC and the RNA was fixed by heating the blot for 30min at 80°C. Probes were generated using "Rediprime™ II random prime labelling system" kit (Amersham) in the presence of (P<sup>32</sup>) dCTP. Hybridisation was carried out at 65°C for 24 h in hybridisation solution (5X SSC, 40µg/ml Denatured Salmon Sperm DNA, 5% [w/v] SDS, and 5× Denhardt's solution). Then, the membrane was washed twice at 65°C using 2X SSC, 0.1% SDS [w/v] and 1X SSC, 0.1% SDS [w/v] during 30 minutes each.

25 5' RACE PCR

The cDNA insert of clone A5-812 was found to contain introns. Therefore, to confirm the coding sequence of this protein, it was necessary to isolate a new cDNA containing the complete coding sequence. This was accomplished by using the SMART<sup>TM</sup> RACE cDNA amplification Kit (Clontech). The first strand cDNA used for the 5' RACE was made as  
5 already described for the cDNA libraries above. A gene specific primer rAP2 (5' CATATAATATTAAAAGCACCAACCATAA 3') was designed - this sequence is situated 92 pb from the poly (A) tail of A5-812 clone. This specific primer was then used with the Universal Primer Mix (UPM) in the CLONTECH kit in a PCR reaction under the following conditions; 2.5 µl of first strand cDNA product, 5 µl of 10X Advantage 2 PCR Buffer  
10 (CLONTECH), 1 µl of dNTP Mix (10 mM), 1 µl of 50X Advantage 2 Polymerase Mix (CLONTECH), 5 µl of "Universal Primer A Mix" (10X) (CLONTECH), 1 µl of rAP2 (10 µM) and sterile water was added to a final volume of 50 µl. PCR cycling conditions were 20 cycles of 30 sec at 94°C, 30 sec at 68°C and 3 min at 72°C, followed by a final extension reaction for 5 min at 72°C. A fragment of about 1700 pb was obtained, excised from the gel  
15 using "CONCERT<sup>TM</sup> Rapid Gel Extraction kit" (GibcoBRL). The isolated fragment was cloned in the pCR 4-TOPO vector and transformed into Escherichia coli using the Topo-TA cloning kit (Invitrogen). The plasmid obtained was then purified using a plasmid extraction kit (QIAfilter Plasmid Midi Kit, Qiagen, France) and the insert of this plasmid was double strand sequenced.

20 The DNA of clone A5-442 (AP1) was found to lack the 5' region of the cDNA. To isolate this region a 5' RACE was performed using the SMART<sup>TM</sup> RACE cDNA amplification Kit (Clontech). A sequence specific primer rAP1 (5'-

25 TGGAGTCACAAGATGTCTCGACGAACTG-3') situated at 396 pb from the poly (A) tail was designed. This specific primer was then used with the Universal Primer Mix (UPM) in the CLONTECH kit in a PCR reaction under the following conditions; 2.5 µl of first strand cDNA, 5 µl of 10X Advantage 2 PCR Buffer (CLONTECH), 1 µl of dNTP Mix (10 mM), 1 µl of 50X Advantage 2 Polymerase Mix (CLONTECH), 5 µl of "Universal Primer A Mix" (10X) (CLONTECH), 1 µl of rAP1, and sterile water was added to a final volume of 50 µl.

30 PCR cycling conditions were 20 cycles of 30 sec at 94°C, 30 sec at 68°C and 3 min at 72°C, followed by a final extension reaction for 5 min at 72°C. A fragment of about 2,000 bp was obtained, excised from the gel using "CONCERT<sup>TM</sup> Rapid Gel Extraction kit" (GibcoBRL). The isolated fragment was cloned in the pCR 4-TOPO vector and transformed into

Escherichia coli using the Topo-TA cloning kit (Invitrogen). The plasmid obtained was then purified using a plasmid extraction kit (QIAfilter Plasmid Midi Kit, Qiagen, France) and the insert of this plasmid was double strand sequenced.

- 5 The Northern blot analysis shown in Figure 1 demonstrates that the coffee cysteine proteinase gene CcCP-1 gene is expressed in the *C. arabica* coffee cherry at all the stages tested, with yellow cherries exhibiting slightly higher levels of expression than the other stages. No expression was detected for this gene in the root, stem or leaves of *C. arabica*.
- 10 The Northern blot analysis shown in Figure 2 demonstrates that the coffee cysteine proteinase inhibitor gene CcCPI-1 gene is expressed in the *C. arabica* coffee cherry at all stages tested. However, in contrast to the expression seen for the cysteine proteinase CcCP-1, CcCPI-1 exhibits higher expression in the two early stages of coffee cherry development (small green and large green), and this gene is expressed at lower levels in the two later stages of cherry
- 15 development. This expression pattern is consistent with the present hypothesis that the cysteine proteinase inhibitor protein (CcCPI-1) controls the activity level of the cysteine proteinase CcCP-1 in the coffee cherry. A controlling protein such as the cysteine proteinase inhibitor protein can be expected to be expressed earlier than its target protein if it is necessary to control the level of activity of its target protein continuously from the time that
- 20 the target protein is expressed. No expression was detected for this gene in the root, stem or leaves of *C. arabica*. It is noted that the similarity of the expression patterns for CcCP-1 and CcCPI-1 are consistent with the present hypothesis that these proteins interact functionally.
- 25 The Northern blot analysis shown in Figure 3 demonstrates that the coffee cysteine proteinase inhibitor gene CcCPI-1 gene is expressed differently in the cherries of *C. canephora* (robusta) versus the cherries of *C. arabica*. First, the data of Figure 3 shows that the CcCPI-1 gene is expressed slightly earlier in *C. arabica*. Secondly, and more importantly, the CcCPI-1 gene is expressed in significantly higher levels in the *C. canephora* cherries. This difference in expression probably affects the level of the cysteine proteinase activity found in *C. arabica*
- 30 versus *C. canephora* cherries. Because this class of protein is widely associated with insect resistance in plants, it is also likely that the high expression of the CcCPI-1 gene in *C. canephora* contributes to the higher disease resistance often seen for robusta varieties versus arabica varieties.

The Northern blot analysis shown in Figure 4 demonstrates that the coffee aspartic proteinase CcAP-2 gene is expressed in both the grain and the pericarp of the *C. arabica* coffee cherry at all cherry development stages tested. The CcAP-2 gene also has a relatively high expression in roots. When the film is exposed longer, CcAP-2 expression was also detected in the tissues of *C. arabica* stems, leaf, and flowers.

*Overexpression and under-expression of the CcCP-1, CcAP-1 and CcAP-2 proteinase gene sequences and the CcCPI-1 proteinase inhibitor in coffee seeds.*

It is expected that the major storage protein profile and the amino acid/peptide profile can be changed in the mature coffee grain by altering, either up or down, the expression of one or more of the genes disclosed herein.

Methods for the overexpression of a gene of interest are well known in the art. Such methods consist of creating a chimeric gene of three major components, 1) a promoter sequence at the 5' end of the gene, preferably in the current application a seed specific promoter such the coffee seed specific promoter described in Marrachini et al. 1999 (Marraccini *et al* 1999 Molecular cloning of the complete 11S seed storage protein gene of *Coffea arabica* and promoter analysis in transgenic tobacco plants, Plant Physiol. Biochem. Vol 37, 273-282, and WO 99/02688), 2) the entire coding sequence of the gene to be expressed, and 3) a 3' control region such as the 3' region from the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. Then, the chimeric gene can be cloned into an *Agrobacterium tumefaciens* transformation vector, and this vector can be transformed into an *Agrobacterium tumefaciens* strain for use in coffee transformation as described by Leroy *et al* 2000, (Leroy *et al* 2000 Genetically modified coffee plants expressing the *Bacillus thuringiensis cry1Ac* gene for resistance to leaf minor. Plant Cell Reports 2000, 19, 382-389). Plants with stable transformation inserts can then be screened for those which overexpress the specific genes used in the transformation experiment specifically in mature seeds using methods such as detection of gene overexpression or protein activity overexpression versus seeds from mock transformed plants.

It is well known in the art that the expression of known gene sequences can be reduced or completely blocked by antisense suppression and of gene expression using nucleic acid fragments representing less than the entire coding region of a gene, and by nucleic acids that do not share 100% sequence identity with the gene to be suppressed. In this case, the

sequences chosen for the particular antisense suppression or cosuppression experiment will replace the full length gene in the chimeric gene construction scheme presented above. The resulting antisense suppression or cosuppression chimeric constructions are again cloned into an *Agrobacterium tumefaciens* transformation vector, and transformed into *Agrobacterium tumefaciens* strain for use in coffee transformation as described above. Plants with stable transformation inserts can then be screened for those with reduced expression of the specific gene sequences used in the seeds of the transformed plants. The reduced expression can be detected by techniques such as northern blotting ;semi quantitative RT-PCR, and/or quantitative RT-PCR.

Another method for reducing, or eliminating, the expression of a gene in plants is to use the small portions of the gene sequences disclosed herein to produce RNA silencing via using RNAi (Hannon, G.J., 2002, Nature, Vol 418, 244-251; Tang *et al*, 2003, Genes Dev, Vol 17, 49-63). In this approach, small regions of one or more of the sequences disclosed herein are cloned into an *Agrobacterium tumefaciens* transformation vector as described above which has a seed specific promoter and an appropriate 3' regulatory region. This new inserted sequence for RNAi should be constructed so that the RNA produced forms an RNA structure *in vivo* which result in the production of small double stranded RNA in the transformed cells and whereby these small double stranded RNA sequences trigger the degradation of the homologous mRNA in these transformed cells.

*Screening for naturally occurring variations in the CcCP-1, CcAP-1, CcAP-2, CcCPI-1 genes and creating new mutations in these genes.*

The sequences disclosed herein can be used to screen natural populations for allelic variants in these genes. This can be accomplished by using the CcCP-1, CcAP-2, CcCPI-1 sequences as probes in a search for naturally occurring RFLP's (restriction fragment length polymorphisms) in genomic DNA from different coffee plant varieties. A more powerful method to find allelic variants is to use the mutation screening technology associated with the TILLING method (Till, B.J., *et al* 2003 Large scale discovery of induced point mutations with high-thruput TILLING. Genome Research Vol 13, 524-530). In this case, once a specific gene sequence has been isolated and cloned, such as CcCP-1, CcAP-2, CcCPI-1 sequences herein, the mutation screening technique associated with the TILLING method can be used to identify sequence variants between the cloned sequence and the corresponding cDNA or genomic sequence in different varieties. Using PCR primer pairs coding for DNA



segments of 700-1100 base pairs, the known cloned gene can be scanned for naturally occurring sequence variations in different varieties. In the ideal situation, one or more sequence variants could also be correlated with a particular phenotypic variation thereby identifying a genetic marker for this phenotypic variant.

5

Additionally, using the sequences disclosed herein for CcCP-1, CcAP-2 and CcCPI-1, application of the full TILLING method can be used to create and detect new mutants in these genes and thus produce plants containing these specific mutants. For example, using the full TILLING method, coffee plants could be created which have specific mutations, such

10

as a missense mutation in the coding sequence which inactivates the gene target of interest.

2005 JUN 18

**CLAIMS:**

1. An isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide  
5 having cysteine proteinase activity, wherein the amino acid sequence of the polypeptide  
and the amino acid sequence of SEQ ID No. 2 have at least 70%, preferably at least 80%,  
sequence identity based on the ClustalW alignment method; or the complement of the  
nucleotide sequence, wherein the complement contains the same number of nucleotides as  
10 the nucleotide sequence, and the complement and the nucleotide sequence are 100%  
complementary.
2. The polynucleotide of Claim 1, wherein the amino acid sequence of the polypeptide and  
the amino acid sequence of SEQ ID No. 2 have at least 85%, preferably at least 90%,  
optionally at least 95%, sequence identity based on the ClustalW alignment method.  
15
3. The polynucleotide of Claim 1, wherein the nucleotide sequence comprises the nucleotide  
sequence of SEQ ID No. 1.
4. The polynucleotide of Claim 1, wherein the polypeptide comprises the amino acid  
20 sequence of SEQ ID No. 2.
5. An isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide  
having cysteine proteinase inhibitor activity, wherein the amino acid sequence of the  
polypeptide and the amino acid sequence of SEQ ID No. 4 have at least 70%, preferably  
25 at least 80%, sequence identity based on the ClustalW alignment method; or the  
complement of the nucleotide sequence, wherein the complement contains the same  
number of nucleotides as the nucleotide sequence, and the complement and the nucleotide  
sequence are 100% complementary.
- 30 6. The polynucleotide of Claim 5, wherein the amino acid sequence of the polypeptide and  
the amino acid sequence of SEQ ID No. 4 have at least 85%, preferably at least 90%,  
optionally at least 95%, sequence identity based on the ClustalW alignment method.

7. The polynucleotide of Claim 5, wherein the nucleotide sequence comprises the nucleotide sequence of SEQ ID No. 3.
8. The polynucleotide of Claim 5, wherein the polypeptide comprises the amino acid sequence of SEQ ID No. 4.
9. An isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide having aspartic endoproteinase activity, wherein the amino acid sequence of the polypeptide and the amino acid sequence selected from SEQ ID No. 6 or 8, preferably SEQ ID No. 8, have at least 75%, preferably at least 80%, sequence identity based on the ClustalW alignment method, or the complement of the nucleotide sequence, wherein the complement contains the same number of nucleotides as the nucleotide sequence, and the complement and the nucleotide sequence are 100% complementary.
10. The polynucleotide of Claim 9, wherein the amino acid sequence of the polypeptide and the amino acid sequence selected from SEQ ID No. 6 or 8, preferably SEQ ID No. 8, have at least 85%, preferably at least 90%, optionally at least 95%, sequence identity based on the ClustalW alignment method.
11. The polynucleotide of Claim 9, wherein the nucleotide sequence comprises the nucleotide sequence of SEQ ID No. 5 or 7, preferably SEQ ID No. 7.
12. The polynucleotide of Claim 9, wherein the polypeptide comprises the amino acid sequence of SEQ ID No. 6 or 8, preferably SEQ ID No. 8.
13. A vector comprising the polynucleotide of any one of Claims 1 to 12.
14. A recombinant DNA construct comprising the polynucleotide of any one of Claims 1 to 12 operably linked to a regulatory sequence.
15. A method for transforming a cell comprising transforming a cell with the polynucleotide of any one of Claims 1 to 12.
16. A cell comprising the recombinant DNA construct of Claim 14.


17. The cell of Claim 16, which is a prokaryotic cell, an eukaryotic cell or a plant cell, preferably a coffee cell.
- 5 18. A transgenic plant comprising the cell of Claim 16 or 17.
19. A method for modulating coffee flavour precursor levels in green coffee grains, the method comprising introducing into the coffee plant the recombinant DNA construct of Claim 14.

## **Abstract**

### **Modulation of Coffee Flavour Precursor Levels in Green Coffee Grains**

- 5    The present invention relates to isolated polynucleotides encoding cysteine proteinases; cysteine proteinase inhibitors; and aspartic endoproteinases.


The invention also relates to a transformed host cell, preferably a plant cell, in which over- or under- expression of these polynucleotides result in altered levels of coffee flavour precursor  
10    levels, specifically, amino group-containing molecules such as amino acids, peptides and proteins, in green coffee grains.



10

15


**Figure 1:** Northern blot analysis of the expression of the cysteine proteinase (CcCP1) gene in different tissues of *Coffea arabica*.



20

**Figure 2:** Northern blot analysis of the expression of the cysteine proteinase inhibitor (CcCPI-1) gene in different tissues of *Coffea arabica*.


5



10 **Figure 3:** Northern blot analysis of the expression of the cysteine proteinase inhibitor gene (CcCPI-1) at different cherry development stages for *Coffea arabica* (ARA) and *Coffea canephora* (ROB).

15

20



25 **Figure 4:** Northern blot analysis of the expression of the aspartic proteinase 2 (CcAP-2) gene in different tissues of *Coffea arabica*.

## SEQUENCE LISTING

5

### SEQUENCE LISTING

10 <110> Tours Nestlé Research Center

15 <120> Implication Of Proteinase And Proteinase Inhibitor In Coffee  
Flavour

20 <130> Patent Proteinase and Proteinase Inhibitor Coffee

<160> 8

25

<170> PatentIn version 3.1

30

<210> 1

<211> 1543

35

<212> DNA

<213> Coffea canephora

40

<220>

<221> mRNA

45

<222> (1)..(1543)

<223>

50

<220>

<221> CDS

55

<222> (122)..(1315)

<223>

60

<400> 1

aagcagtggg aacaacgcag agtacgcggg ggacactcct ccccgttcca ttccagacca

60



	gggtccaaaa	ccaccgtcca	agagaggagc	agactgcaga	gtgatacata	caggcacaaa	120											
5	g	atg	atg	atg	aca	agc	gga	ggt	ctg	atg	cta	acc	tgc	act	ctg	gct	att	169
	Met	Met	Met	Met	Thr	Ser	Gly	Gly	Leu	Met	Leu	Thr	Cys	Thr	Leu	Ala	Ile	
	1				5					10					15			
10	acc	ctc	tta	tcc	tgc	gca	ctc	atc	tct	tca	acc	act	ttc	caa	cat	gaa		217
	Thr	Leu	Leu	Ser	Cys	Ala	Leu	Ile	Ser	Ser	Thr	Thr	Phe	Gln	His	Glu		
				20				25						30				
15	att	cag	tat	cga	gta	caa	gac	ccg	tta	atg	ata	cgc	caa	gtc	acc	gac		265
	Ile	Gln	Tyr	Arg	Val	Gln	Asp	Pro	Leu	Met	Ile	Arg	Gln	Val	Thr	Asp		
			35					40					45					
20	aat	cac	cac	cac	cgc	cac	cac	cca	ggt	agg	tct	tct	gca	aac	cat	cgt		313
	Asn	His	His	His	Arg	His	His	Pro	Gly	Arg	Ser	Ser	Ala	Asn	His	Arg		
		50					55					60						
25	cta	ctg	ggc	acc	acc	aca	gag	gtt	cac	ttc	aag	tcc	ttc	gtg	gag	gag		361
	Leu	Leu	Gly	Thr	Thr	Thr	Glu	Val	His	Phe	Lys	Ser	Phe	Val	Glu	Glu		
	65					70					75					80		
30	tac	gag	aaa	act	tac	tct	acg	cac	gag	gag	tac	gtg	cac	cgc	ctg	ggg		409
	Tyr	Glu	Lys	Thr	Tyr	Ser	Thr	His	Glu	Glu	Tyr	Val	His	Arg	Leu	Gly		
				85						90					95			
35	att	ttc	gcc	aag	aac	ctc	atc	aag	gcc	gcg	gag	cac	cag	gcc	atg	gac		457
	Ile	Phe	Ala	Lys	Asn	Leu	Ile	Lys	Ala	Ala	Glu	His	Gln	Ala	Met	Asp		
				100					105					110				
40	ccc	tcc	gca	atc	cac	ggc	gtc	acc	cag	ttc	tct	gat	ctc	acc	gag	gag		505
	Pro	Ser	Ala	Ile	His	Gly	Val	Thr	Gln	Phe	Ser	Asp	Leu	Thr	Glu	Glu		
			115					120					125					
45	gag	ttt	gag	gct	acg	tac	atg	ggc	ctt	aaa	ggt	ggc	gct	gga	gtt	ggt		553
	Glu	Phe	Glu	Ala	Thr	Tyr	Met	Gly	Leu	Lys	Gly	Gly	Ala	Gly	Val	Gly		
		130					135				140							
50	ggg	acc	acc	cag	ctg	ggg	aaa	gat	gat	ggg	gat	gag	agt	gca	gca	gag		601
	Gly	Thr	Thr	Gln	Leu	Gly	Lys	Asp	Asp	Gly	Asp	Glu	Ser	Ala	Ala	Glu		
	145				150				155							160		
55	gtg	atg	atg	gat	gta	tct	gat	ttg	ccg	gag	agt	ttt	gat	tgg	aga	gaa		649
	Val	Met	Met	Asp	Val	Ser	Asp	Leu	Pro	Glu	Ser	Phe	Asp	Trp	Arg	Glu		
				165					170					175				
60	aaa	ggt	gct	gtg	acc	gaa	gtg	aag	acg	cag	gga	aga	tgt	gga	tcg	tgt		697
	Lys	Gly	Ala	Val	Thr	Glu	Val	Lys	Thr	Gln	Gly	Arg	Cys	Gly	Ser	Cys		
				180				185					190					
65	tgg	gct	ttt	agt	aca	act	gga	gct	att	gaa	gga	gct	aat	ttc	att	gca		745
	Trp	Ala	Phe	Ser	Thr	Thr	Gly	Ala	Ile	Glu	Gly	Ala	Asn	Phe	Ile	Ala		
			195				200						205					
70	act	ggc	aag	ctt	ctc	agc	cta	agt	gaa	cag	cag	ctt	gtg	gat	tgt	gat		793
	Thr	Gly	Lys	Leu	Leu	Ser	Leu	Ser	Glu	Gln	Gln	Leu	Val	Asp	Cys	Asp		
		210					215					220						
75	cat	atg	tgt	gat	tta	aaa	gaa	aaa	gat	gac	tgt	gat	gat	gga	tgc	tcc		841
	His	Met	Cys	Asp	Leu	Lys	Glu	Lys	Asp	Asp	Cys	Asp	Asp	Gly	Cys	Ser		
	225					230					235					240		
80	gga	ggg	cta	atg	aca	act	gct	ttc	aac	tac	ttg	ata	gag	gca	gga	ggt		889
	Gly	Gly	Leu	Met	Thr	Thr	Ala	Phe	Asn	Tyr	Leu	Ile	Glu	Ala	Gly	Gly		
				245				250						255				
85	ata	gag	gag	gag	gta	acc	tat	ccc	tac	act	ggg	aaa	cgc	gga	gaa	tgc		937
	Ile	Glu	Glu	Glu	Val	Thr	Tyr	Pro	Tyr	Thr	Gly	Lys	Arg	Gly	Glu	Cys		
				260				265						270				

5	aaa ttc aat cct gag aaa gtt gcg gtg aaa gtg cgg aat ttc gca aaa Lys Phe Asn Pro Glu Lys Val Ala Val Lys Val Arg Asn Phe Ala Lys	985
10	atc cct gag gat gag agt caa att gct gcc aat gta gtg cat aat ggc Ile Pro Glu Asp Glu Ser Gln Ile Ala Ala Asn Val Val His Asn Gly	1033
15	ccg ctt gct att gga ttg aat gcg gta ttc atg caa act tac atc ggg Pro Leu Ala Ile Gly Leu Asn Ala Val Phe Met Gln Thr Tyr Ile Gly	1081
20	ggt gtg tca tgt cct ctt att tgt gac aaa aag agg atc aac cat ggt Gly Val Ser Cys Pro Leu Ile Cys Asp Lys Lys Arg Ile Asn His Gly	1129
25	gtt ctt ctt gtg ggc tat ggt tct aga ggc ttc tca atc ctt agg ctt Val Leu Leu Val Gly Tyr Gly Ser Arg Gly Phe Ser Ile Leu Arg Leu	1177
30	ggc tac aag cca tac tgg att atc aag aac tca tgg ggg aag cgt tgg Gly Tyr Lys Pro Tyr Trp Ile Ile Lys Asn Ser Trp Gly Lys Arg Trp	1225
35	ggc gaa cat ggt tgc tac cgg ctt tgt cga ggg cac aac atg tgt gga Gly Glu His Gly Cys Tyr Arg Leu Cys Arg Gly His Asn Met Cys Gly	1273
40	atg agc aca atg gtt tca gct gtg gtg aca cag acc tct tga Met Ser Thr Met Val Ser Ala Val Val Thr Gln Thr Ser	1315
45	taccaaaaca tctctgctct tcagaggttg tatacaaggt ggtttgctct tggaagatct	1375
50	tatcatgttt tcgaaatatt taggtttgta taatatgaag ggtagagagt aataagaacc	1435
55	aaacaaaagt tcaggcctgt ttctgatagg aatggaatat gatcggagtc atttggtact	1495
60	ggatcacaaa aaaaaatcca aaaaaaaaaa aaaaaaaaaa aaaaaaaa	1543
65	<210> 2	
70	<211> 397	
75	<212> PRT	
80	<213> Coffea canephora	
85	<400> 2	
90	Met Met Met Thr Ser Gly Gly Leu Met Leu Thr Cys Thr Leu Ala Ile 1 5 10 15	
95	Thr Leu Leu Ser Cys Ala Leu Ile Ser Ser Thr Thr Phe Gln His Glu 20 25 30	
100	Ile Gln Tyr Arg Val Gln Asp Pro Leu Met Ile Arg Gln Val Thr Asp 35 40 45	
105	Asn His His His Arg His His Pro Gly Arg Ser Ser Ala Asn His Arg 50 55 60	

5 Leu 65 Leu Gly Thr Thr Thr 70 Glu Val His Phe Lys 75 Ser Phe Val Glu 80 Glu  
 Tyr Glu Lys Thr Tyr 85 Ser Thr His Glu 90 Glu Tyr Val His Arg Leu 95 Gly  
 10 Ile Phe Ala Lys 100 Asn Leu Ile Lys Ala 105 Ala Glu His Gln Ala 110 Met Asp  
 15 Pro Ser Ala 115 Ile His Gly Val Thr 120 Gln Phe Ser Asp Leu 125 Thr Glu Glu  
 20 Glu Phe 130 Glu Ala Thr Tyr Met 135 Gly Leu Lys Gly 140 Gly Ala Gly Val Gly  
 25 Gly 145 Thr Thr Gln Leu Gly 150 Lys Asp Asp Gly 155 Asp Glu Ser Ala Ala Glu 160  
 30 Val Met Met Asp Val 165 Ser Asp Leu Pro Glu 170 Ser Phe Asp Trp Arg 175 Glu  
 Lys Gly Ala Val 180 Thr Glu Val Lys Thr 185 Gln Gly Arg Cys Gly 190 Ser Cys  
 35 Trp Ala Phe 195 Ser Thr Thr Gly Ala 200 Ile Glu Gly Ala Asn 205 Phe Ile Ala  
 40 Thr Gly 210 Lys Leu Leu Ser Leu 215 Ser Glu Gln Gln Leu 220 Val Asp Cys Asp  
 45 His 225 Met Cys Asp Leu Lys 230 Glu Lys Asp Asp Cys 235 Asp Asp Gly Cys Ser 240  
 Gly Gly Leu Met Thr 245 Thr Ala Phe Asn Tyr 250 Leu Ile Glu Ala Gly 255 Gly  
 50 Ile Glu Glu Glu 260 Val Thr Tyr Pro Tyr 265 Thr Gly Lys Arg Gly 270 Glu Cys  
 55 Lys Phe Asn 275 Pro Glu Lys Val Ala 280 Val Lys Val Arg Asn 285 Phe Ala Lys  
 60 Ile Pro 290 Glu Asp Glu Ser Gln 295 Ile Ala Ala Asn Val 300 Val His Asn Gly  
 Pro 305 Leu Ala Ile Gly Leu 310 Asn Ala Val Phe Met 315 Gln Thr Tyr Ile Gly 320  
 65 Gly Val Ser Cys Pro 325 Leu Ile Cys Asp Lys 330 Lys Arg Ile Asn His 335 Gly

	Val	Leu	Leu	Val	Gly	Tyr	Gly	Ser	Arg	Gly	Phe	Ser	Ile	Leu	Arg	Leu	
				340					345					350			
5	Gly	Tyr	Lys	Pro	Tyr	Trp	Ile	Ile	Lys	Asn	Ser	Trp	Gly	Lys	Arg	Trp	
			355					360					365				
10	Gly	Glu	His	Gly	Cys	Tyr	Arg	Leu	Cys	Arg	Gly	His	Asn	Met	Cys	Gly	
		370					375					380					
15	Met	Ser	Thr	Met	Val	Ser	Ala	Val	Val	Thr	Gln	Thr	Ser				
	385					390					395						
	<210>	3															
20	<211>	726															
	<212>	DNA															
	<213>	Coffea canephora															
25	<220>																
30	<221>	mRNA															
	<222>	(1)..(726)															
	<223>																
35	<220>																
40	<221>	CDS															
	<222>	(79)..(498)															
	<223>																
45	<400>	3															
	ggcgc	aaacaa	acattg	aaag	aaaatc	aaga	acccaaaaa	accccaca	ag	aaaaaa	agaa						60
50	aaaga	agaag	aaaagcca	atg	gca	aaa	cca	tcg	tca	tct	cta	ctc	aca	ctt			111
				Met	Ala	Lys	Pro	Ser	Ser	Ser	Leu	Leu	Thr	Leu			
				1				5					10				
55	cct	tcc	ttt	ctt	ctg	atc	ttt	ttc	att	ctt	gca	cta	ttt	tcc	acc	acc	159
	Pro	Ser	Phe	Leu	Leu	Ile	Phe	Phe	Ile	Leu	Ala	Leu	Phe	Ser	Thr	Thr	
				15					20					25			
60	ctc	caa	gtt	aat	gcc	ttg	gga	agg	aaa	gtg	gga	gca	agg	gag	aag	att	207
	Leu	Gln	Val	Asn	Ala	Leu	Gly	Arg	Lys	Val	Gly	Ala	Arg	Glu	Lys	Ile	
			30				35					40					
65	gag	gat	gtg	aag	agc	aac	aaa	gaa	gtt	caa	gaa	ctt	ggg	gaa	tat	tgt	255
	Glu	Asp	Val	Lys	Ser	Asn	Lys	Glu	Val	Gln	Glu	Leu	Gly	Glu	Tyr	Cys	
		45					50					55					
	gtt	tct	gag	tac	aac	aag	agt	ttg	cgg	aag	aag	aac	aac	gaa	agt	ggt	303
	Val	Ser	Glu	Tyr	Asn	Lys	Ser	Leu	Arg	Lys	Lys	Asn	Asn	Glu	Ser	Gly	
	60					65					70					75	

gct cct ata atc ttc aca tct gtg gtg gag gct gag aag cag gtg gtt 351  
 Ala Pro Ile Ile Phe Thr Ser Val Val Glu Ala Glu Lys Gln Val Val  
 80 85 90

5 gct ggg atc aaa tat tat ctc aag att aag gcc acc act tct tct ggg 399  
 Ala Gly Ile Lys Tyr Tyr Leu Lys Ile Lys Ala Thr Thr Ser Ser Gly  
 95 100 105

10 gtt ccc aag gtt tac gat gcc att gtg gtg gtt cgg cct tgg gtt cat 447  
 Val Pro Lys Val Tyr Asp Ala Ile Val Val Val Arg Pro Trp Val His  
 110 115 120

15 act aag cca agg cag ttg ctc aac ttc tcc cct tcc cct gcc act aaa 495  
 Thr Lys Pro Arg Gln Leu Leu Asn Phe Ser Pro Ser Pro Ala Thr Lys  
 125 130 135

tga agaagaaaat gttgaaaaag ttggaactgt ttgggagatc taatctgatg 548

20 attattagta cctttcagtg caaattctct ttgctgttaa gtgttcgggtt tttttttttt 608

ccctgtgtct atttatgacc gtgggtcatga tgatatggtg tatgatccag taataattaa 668

aatctgttgc ataaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaa 726

25 <210> 4

<211> 139

30 <212> PRT

<213> Coffea canephora

35 <400> 4

40 Met Ala Lys Pro Ser Ser Ser Leu Leu Thr Leu Pro Ser Phe Leu Leu  
 1 5 10 15

Ile Phe Phe Ile Leu Ala Leu Phe Ser Thr Thr Leu Gln Val Asn Ala  
 20 25 30

45 Leu Gly Arg Lys Val Gly Ala Arg Glu Lys Ile Glu Asp Val Lys Ser  
 35 40 45

50 Asn Lys Glu Val Gln Glu Leu Gly Glu Tyr Cys Val Ser Glu Tyr Asn  
 50 55 60

55 Lys Ser Leu Arg Lys Lys Asn Asn Glu Ser Gly Ala Pro Ile Ile Phe  
 65 70 75 80

60 Thr Ser Val Val Glu Ala Glu Lys Gln Val Val Ala Gly Ile Lys Tyr  
 85 90 95

Tyr Leu Lys Ile Lys Ala Thr Thr Ser Ser Gly Val Pro Lys Val Tyr  
 100 105 110

65 Asp Ala Ile Val Val Val Arg Pro Trp Val His Thr Lys Pro Arg Gln  
 115 120 125

Leu Leu Asn Phe Ser Pro Ser Pro Ala Thr Lys  
130 135

5 <210> 5  
<211> 2282  
10 <212> DNA  
<213> Coffea canephora  
15 <220>  
<221> mRNA  
20 <222> (1)..(2282)  
<223>  
25 <220>  
<221> CDS  
30 <222> (439)..(1731)  
<223>  
35 <400> 5  
actcactata ctttgcattc tcttcacccat tctccctcaa aactccctcc aacattcttt 60  
tccttggttt tttcatctat ccctcctata aaaatcgatt attttggttct tttacctctt 120  
40 aaaaatccat tcttggaatt catttatcca tatacaccat acttggtgcat gtcccttttg 180  
gttgttttgc ttttggtgata agtaattggt gggtttattgg tttttcatga tggctccgga 240  
45 tctaagaaga aatggggtcgg tagtagcttt agccctgtta gtctctctgg ttgttaatgg 300  
tggtattttt gatgtagaag gtaacaataa tgtgggtttt gaggtggaac ataaatttaa 360  
agggagaagg aatgagaatg gaggaagagg gtctttttga cttcactcaa ggctcatgat 420  
50 tcccaccgcc atggcaga atg ctt gca gcc ctt gac atg cct ttg ggt ggc 471  
Met Leu Ala Ala Leu Asp Met Pro Leu Gly Gly  
1 5 10  
55 aat ggt tcc cct aca gat gca gcg ctc tat ttc act aag ctt tcg att 519  
Asn Gly Ser Pro Thr Asp Ala Ala Leu Tyr Phe Thr Lys Leu Ser Ile  
15 20 25  
60 ggg act cct cct cag gat tat tat gtg caa gtg gat aca gga agt gac 567  
Gly Thr Pro Pro Gln Asp Tyr Tyr Val Gln Val Asp Thr Gly Ser Asp  
30 35 40  
65 att ctc tgg gta aac tgt gct ggt tgt gtc aga tgc ccc aag aaa agc 615  
Ile Leu Trp Val Asn Cys Ala Gly Cys Val Arg Cys Pro Lys Lys Ser  
45 50 55  
agt ctt ggt att gac ttg act cta tat gac atg aaa gcc tcc agc acc 663  
Ser Leu Gly Ile Asp Leu Thr Leu Tyr Asp Met Lys Ala Ser Ser Thr  
60 65 70 75

	ggg aga ctt gtt act tgt gat caa gac ttt tgc ttg tct gca ttc aat	711
	Gly Arg Leu Val Thr 80 Cys Asp Gln Asp Phe 85 Cys Leu Ser Ala Phe 90 Asn	
5	gcc cca gcc tct gat tgc aag gtt ggt aac ccc tgt gca tat tct gtt	759
	Ala Pro Ala Ser 95 Asp Cys Lys Val 100 Gly Asn Pro Cys Ala Tyr 105 Ser Val	
10	act tac gga gac ggg agc tca acc ggc gga tat ttt gtc aga gac tat	807
	Thr Tyr Gly Asp Gly Ser Ser Thr 115 Gly Gly Tyr Phe Val 120 Arg Asp Tyr	
15	gca aaa ctt aat caa ctg acg gga aat ctt caa acc ata ccc atg aat	855
	Ala Lys 125 Leu Asn Gln Leu Thr 130 Gly Asn Leu Gln Thr 135 Ile Pro Met Asn	
20	ggt agt ata gtg ttt ggg tgt tca tct caa caa tct gga gag cta ggg	903
	Gly 140 Ser Ile Val Phe 145 Gly Cys Ser Ser Gln Gln 150 Ser Gly Glu Leu Gly 155	
25	tca tct act gaa gca gtt gat ggc ata att ggt ttt gga caa gca aat	951
	Ser Ser Thr Glu Ala Val Asp Gly Ile 165 Gly Phe Gly Gln Ala Asn 170	
30	tca tct att att tca cag ctt gct tca gca gga aag gtt aaa aaa ata	999
	Ser Ser Ile Ile Ser Gln Leu Ala Ser Ala Gly Lys Val 185 Lys Lys Ile 175	
35	ttt tca cat tgc ttg gat ggt atc aat gga gga ggc ata ttt gct att	1047
	Phe Ser His 190 Cys Leu Asp Gly Ile 195 Asn Gly Gly Gly Ile 200 Phe Ala Ile	
40	gga caa gta gtg cag cca aaa cta aag aca aca cca ttg gtc cca aat	1095
	Gly Gln Val Val Gln Pro Lys 210 Leu Lys Thr Thr Pro 215 Leu Val Pro Asn 205	
45	gag gca cat tat aat gtt gtt ctg aac gca att gag gtg ggt ggc gac	1143
	Glu Ala His Tyr Asn Val Val Leu Asn Ala Ile 230 Glu Val Gly Gly Asp 235 220	
50	ggt cta aac ctt ccc tca gat gta tta gga ggt gga tct gga agt ggt	1191
	Val Leu Asn Leu Pro 240 Ser Asp Val Leu Gly 245 Gly Gly Ser Gly Ser 250 Gly	
55	aca ata ata gac agt ggt aca acc ttg gct tat ctt cct gat gat gtc	1239
	Thr Ile Ile Asp Ser Gly Thr Thr Leu Ala Tyr Leu Pro 265 Asp Asp Val 255	
60	tat act cca ctt atg gaa aag att acg gca tcc caa tcc aac ttg aaa	1287
	Tyr Thr Pro 270 Leu Met Glu Lys Ile 275 Thr Ala Ser Gln Ser 280 Asn Leu Lys	
65	atc cat att gtt gaa aat cag ttc aag tgc ttt gtc tat agt gga aat	1335
	Ile His 285 Ile Val Glu Asn Gln 290 Phe Lys Cys Phe Val 295 Tyr Ser Gly Asn	
70	ggt gat gat gga ttt cca gtt gta tct ttc cac ttt gag gat tca ctt	1383
	Val Asp Asp Gly Phe 305 Val Val Ser Phe His 310 Phe Glu Asp Ser Leu 315 300	
75	tct ttg aca gtt tat ccc cat gaa tat ctc ttt gat ctt cat gat gat	1431
	Ser Leu Thr Val Tyr 320 Pro His Glu Tyr Leu Phe 325 Asp Leu His Asp Asp 330	
80	caa tgg tgt att ggt tgg cag aat aag ggt atg cag aca aga gat gga	1479
	Gln Trp Cys Ile 335 Gly Trp Gln Asn Lys 340 Gly Met Gln Thr Arg 345 Asp Gly	
85	agg gaa gta act ctt ttg gga gat ctt gta ctc gca aac aag ctt gtt	1527

Arg Glu Val Thr Leu Leu Gly Asp Leu Val Leu Ala Asn Lys Leu Val  
 350 355 360  
 5 tct tat gat ctt gaa aat caa acc att gga tgg gct gaa tat aat tgc 1575  
 Ser Tyr Asp Leu Glu Asn Gln Thr Ile Gly Trp Ala Glu Tyr Asn Cys 375  
 10 tct tct agc atc aaa ttg aga gac gag aag tca gga aac gtg tat gct 1623  
 Ser Ser Ser Ile Lys Leu Arg Asp Glu Lys Ser Gly Asn Val Tyr Ala 395  
 15 gtg ggc tct cat atc att tct tca gct cgc ggc ctg aat gct gga aag 1671  
 Val Gly Ser His Ile Ile Ser Ser Ala Arg Gly Leu Asn Ala Gly Lys 410  
 20 gct cta agg ttc cta ttg tta atc atc aca tca ttg ttg cat gca ctt 1719  
 Ala Leu Arg Phe Leu Leu Leu Ile Ile Thr Ser Leu Leu His Ala Leu 425  
 25 ttg atc cca tga acatttaaaa tcatactagc tgagaaggag gcattatgat 1771  
 Leu Ile Pro 430  
 30 agcgtaccat ggtactcata gtgatcaggc atcttgctga ttctttggac cattataatt 1831  
 tctcatgtgt ttaaagtgaag agtcagttcg tcgagacatc ttgtgactcc ataattctct 1891  
 tgatcaagct gaactctact cacaaaacca tagctaattc ttttgatctc aaaagagaaa 1951  
 taggctctgc aaaaggattt cggagggttga tggtgaacat tcttcttatt tggatgttat 2011  
 tgatacccca gatgattaag gaaagcctat aggaaacaga tgggtgggaag gagtatacat 2071  
 35 tctttctgac tctttggaac ttcctagcgt atacacatat ttcacacgga atgtatctta 2131  
 taattcatct gttctttctg tttattgtca acttgtttca aatgattgga gtagctgcaa 2191  
 taatcaactc ggatgggtggg tcatgcttaa ggctcgtctt gcctcattgt taagacgtga 2251  
 40 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 2282  
 <210> 6  
 45 <211> 430  
 <212> PRT  
 50 <213> Coffea canephora  
 <400> 6  
 55 Met Leu Ala Ala Leu Asp Met Pro Leu Gly Gly Asn Gly Ser Pro Thr  
 1 5 10 15  
 60 Asp Ala Ala Leu Tyr Phe Thr Lys Leu Ser Ile Gly Thr Pro Pro Gln  
 20 25 30  
 65 Asp Tyr Tyr Val Gln Val Asp Thr Gly Ser Asp Ile Leu Trp Val Asn  
 35 40 45  
 Cys Ala Gly Cys Val Arg Cys Pro Lys Lys Ser Ser Leu Gly Ile Asp  
 50 55 60



5 Leu Thr Leu Tyr Asp Met Lys Ala Ser Ser Thr Gly Arg Leu Val Thr  
 65 70 75 80  
 Cys Asp Gln Asp Phe Cys Leu Ser Ala Phe Asn Ala Pro Ala Ser Asp  
 85 90 95  
 10 Cys Lys Val Gly Asn Pro Cys Ala Tyr Ser Val Thr Tyr Gly Asp Gly  
 100 105 110  
 15 Ser Ser Thr Gly Gly Tyr Phe Val Arg Asp Tyr Ala Lys Leu Asn Gln  
 115 120 125  
 20 Leu Thr Gly Asn Leu Gln Thr Ile Pro Met Asn Gly Ser Ile Val Phe  
 130 135 140  
 Gly Cys Ser Ser Gln Gln Ser Gly Glu Leu Gly Ser Ser Thr Glu Ala  
 145 150 155 160  
 25 Val Asp Gly Ile Ile Gly Phe Gly Gln Ala Asn Ser Ser Ile Ile Ser  
 165 170 175  
 30 Gln Leu Ala Ser Ala Gly Lys Val Lys Lys Ile Phe Ser His Cys Leu  
 180 185 190  
 35 Asp Gly Ile Asn Gly Gly Gly Ile Phe Ala Ile Gly Gln Val Val Gln  
 195 200 205  
 40 Pro Lys Leu Lys Thr Thr Pro Leu Val Pro Asn Glu Ala His Tyr Asn  
 210 215 220  
 Val Val Leu Asn Ala Ile Glu Val Gly Gly Asp Val Leu Asn Leu Pro  
 225 230 235 240  
 45 Ser Asp Val Leu Gly Gly Gly Ser Gly Ser Gly Thr Ile Ile Asp Ser  
 245 250 255  
 50 Gly Thr Thr Leu Ala Tyr Leu Pro Asp Asp Val Tyr Thr Pro Leu Met  
 260 265 270  
 55 Glu Lys Ile Thr Ala Ser Gln Ser Asn Leu Lys Ile His Ile Val Glu  
 275 280 285  
 60 Asn Gln Phe Lys Cys Phe Val Tyr Ser Gly Asn Val Asp Asp Gly Phe  
 290 295 300  
 Pro Val Val Ser Phe His Phe Glu Asp Ser Leu Ser Leu Thr Val Tyr  
 305 310 315 320  
 65 Pro His Glu Tyr Leu Phe Asp Leu His Asp Asp Gln Trp Cys Ile Gly  
 325 330 335

Trp Gln Asn Lys Gly Met Gln Thr Arg Asp Gly Arg Glu Val Thr Leu  
 340 345 350  
 5 Leu Gly Asp Leu Val Leu Ala Asn Lys Leu Val Ser Tyr Asp Leu Glu  
 355 360 365  
 10 Asn Gln Thr Ile Gly Trp Ala Glu Tyr Asn Cys Ser Ser Ser Ile Lys  
 370 375 380  
 15 Leu Arg Asp Glu Lys Ser Gly Asn Val Tyr Ala Val Gly Ser His Ile  
 385 390 395 400  
 20 Ile Ser Ser Ala Arg Gly Leu Asn Ala Gly Lys Ala Leu Arg Phe Leu  
 405 410 415  
 25 <210> 7  
 <211> 1819  
 30 <212> DNA  
 <213> Coffea canephora  
 35 <220>  
 <221> mRNA  
 40 <222> (1)..(1732)  
 <223>  
 45 <220>  
 <221> CDS  
 50 <222> (79)..(1602)  
 <223>  
 55 <400> 7  
 cttactgact ctcgtatatt attcaatcta tcttttgagt tttgcaagag cccatcaagc 60  
 atcaaggcat aaccaacg atg gag agg agg tac ctt tgg gca gca ttt gta 111  
 60 Met Glu Arg Arg Tyr Leu Trp Ala Ala Phe Val  
 1 5 10  
 tta ggg gcg att gtg tgt tct cta ttt cct ctt cct tct gaa gga tta 159  
 65 Leu Gly Ala Ile Val Cys Ser Leu Phe Pro Leu Pro Ser Glu Gly Leu  
 15 20 25  
 aag cga att agc ctg aaa aaa aaa ccc tta gat att caa agc ata aga 207  
 30 35 40  
 Lys Arg Ile Ser Leu Lys Lys Pro Leu Asp Ile Gln Ser Ile Arg

		gct	gcc	aaa	tta	gct	cat	ctg	gag	agc	aca	cat	ggc	gct	ggt	agg	aaa	255
		Ala	Ala	Lys	Leu	Ala	His	Leu	Glu	Ser	Thr	His	Gly	Ala	Gly	Arg	Lys	
			45					50					55					
5		gag	atg	gac	aac	aat	tta	ggc	agt	tcc	aat	gag	gac	ata	ttg	cct	tta	303
		Glu	Met	Asp	Asn	Asn	Leu	Gly	Ser	Ser	Asn	Glu	Asp	Ile	Leu	Pro	Leu	
		60					65					70					75	
10		aag	aat	tac	ctg	gat	gcc	cag	tac	tat	gga	gag	att	gga	att	ggt	act	351
		Lys	Asn	Tyr	Leu	Asp	Ala	Gln	Tyr	Tyr	Gly	Glu	Ile	Gly	Ile	Gly	Thr	
						80					85					90		
15		cca	cct	cag	aag	ttc	aca	ggt	ata	ttt	gat	aca	ggc	agt	tcc	aac	ctc	399
		Pro	Pro	Gln	Lys	Phe	Thr	Val	Ile	Phe	Asp	Thr	Gly	Ser	Ser	Asn	Leu	
					95					100					105			
20		tgg	gtg	cca	tcg	gca	aaa	tgt	tac	ttc	tct	att	gcc	tgc	tgg	ctc	cac	447
		Trp	Val	Pro	Ser	Ala	Lys	Cys	Tyr	Phe	Ser	Ile	Ala	Cys	Trp	Leu	His	
				110					115					120				
25		tcc	aag	tac	aag	gca	aag	aag	tca	agt	act	tat	aca	gcc	ata	ggg	aaa	495
		Ser	Lys	Tyr	Lys	Ala	Lys	Lys	Ser	Ser	Thr	Tyr	Thr	Ala	Ile	Gly	Lys	
			125					130					135					
30		tct	tgt	tca	att	cgt	tat	ggg	tct	gga	tca	att	tct	gga	ttc	tcc	agt	543
		Ser	Cys	Ser	Ile	Arg	Tyr	Gly	Ser	Gly	Ser	Ile	Ser	Gly	Phe	Ser	Ser	
		140					145					150					155	
35		cag	gat	aac	ggt	gaa	ggt	ggg	gat	ctt	ggt	gtc	aaa	gat	caa	ggt	ttt	591
		Gln	Asp	Asn	Val	Glu	Val	Gly	Asp	Leu	Val	Val	Lys	Asp	Gln	Val	Phe	
						160					165					170		
40		att	gaa	gct	tca	cga	gaa	gga	agt	ctt	aca	ttt	gta	att	gcc	aag	ttt	639
		Ile	Glu	Ala	Ser	Arg	Glu	Gly	Ser	Leu	Thr	Phe	Val	Ile	Ala	Lys	Phe	
					175					180					185			
45		gac	ggg	ata	ctt	ggc	ctt	gga	ttc	cag	gag	atc	gct	ggt	gat	aac	atg	687
		Asp	Gly	Ile	Leu	Gly	Leu	Gly	Phe	Gln	Glu	Ile	Ala	Val	Asp	Asn	Met	
				190					195					200				
50		gtg	ccg	gtc	tgg	tat	aat	atg	gtg	gac	caa	ggt	ctc	gtg	gat	gag	caa	735
		Val	Pro	Val	Trp	Tyr	Asn	Met	Val	Asp	Gln	Gly	Leu	Val	Asp	Glu	Gln	
			205					210					215					
55		gta	ttc	tct	ttc	tgg	ctt	aac	cgc	gac	cca	aat	gct	gaa	gac	gga	ggt	783
		Val	Phe	Ser	Phe	Trp	Leu	Asn	Arg	Asp	Pro	Asn	Ala	Glu	Asp	Gly	Gly	
		220					225					230					235	
60		gag	ctg	gtc	ttt	ggg	ggg	gta	gat	aca	aat	cac	ttc	aag	gga	aag	cat	831
		Glu	Leu	Val	Phe	Gly	Gly	Val	Asp	Thr	Asn	His	Phe	Lys	Gly	Lys	His	
					240						245					250		
65		aca	tat	ggt	cct	gta	act	cag	aag	gga	tac	tgg	caa	ttt	aaa	atg	gga	879
		Thr	Tyr	Val	Pro	Val	Thr	Gln	Lys	Gly	Tyr	Trp	Gln	Phe	Lys	Met	Gly	
					255					260					265			
70		gat	ttt	ctc	att	ggg	aac	gtc	tca	aca	ggc	ttt	tgt	gaa	gga	ggt	tgt	927
		Asp	Phe	Leu	Ile	Gly	Asn	Val	Ser	Thr	Gly	Phe	Cys	Glu	Gly	Gly	Cys	
				270					275					280				
75		gct	gct	att	gtg	gac	tct	gga	aca	tcg	ttg	ctc	gct	ggt	cca	act	act	975
		Ala	Ala	Ile	Val	Asp	Ser	Gly	Thr	Ser	Leu	Leu	Ala	Gly	Pro	Thr	Thr	
								290					295					
80		gtt	gtg	act	caa	att	aat	cat	gcc	att	gga	gct	gaa	gga	gta	gtt	agc	1023
		Val	Val	Thr	Gln	Ile	Asn	His	Ala	Ile	Gly	Ala	Glu	Gly	Val	Val	Ser	
		300					305					310					315	
		act	gaa	tgt	aaa	gaa	att	gtt	tca	cag	tat	ggt	gaa	ctg	att	tgg	gat	1071

	Thr	Glu	Cys	Lys	Glu 320	Ile	Val	Ser	Gln	Tyr 325	Gly	Glu	Leu	Ile	Trp 330	Asp	
5	ctc Leu	ctc Leu	gta Val	tca Ser 335	ggg Gly	gta Val	cta Leu	ccc Pro	gac Asp 340	aga Arg	gtt Val	tgt Cys	aaa Lys	caa Gln 345	gct Ala	ggt Gly	1119
10	tta Leu	tgt Cys	ccc Pro 350	ctt Leu	cgt Arg	ggt Gly	gct Ala	cag Gln 355	cat His	gag Glu	aat Asn	gct Ala	tat Tyr 360	atc Ile	aag Lys	tca Ser	1167
15	gtc Val	gtc Val 365	gac Asp	gag Glu	gag Glu	aac Asn	aag Lys 370	gag Glu	gaa Glu	gct Ala	tct Ser	gtt Val 375	ggt Gly	gaa Glu	tcc Ser	ccg Pro	1215
20	atg Met 380	tgt Cys	act Thr	gct Ala	tgt Cys	gaa Glu 385	atg Met	gct Ala	gtt Val	gtt Val	tgg Trp 390	atg Met	caa Gln	aac Asn	cag Gln	ctg Leu 395	1263
25	aaa Lys	cag Gln	cag Gln	gga Gly	act Thr 400	aag Lys	gag Glu	aaa Lys	gtg Val	ctt Leu 405	gca Ala	tat Tyr	gtg Val	aat Asn	cag Gln 410	ctt Leu	1311
30	tgt Cys	gaa Glu	agc Ser	ata Ile 415	cca Pro	agt Ser	ccc Pro	atg Met	gga Gly 420	gaa Glu	tcc Ser	atc Ile	att Ile	gac Asp 425	tgc Cys	aac Asn	1359
35	agt Ser	tta Leu	tcc Ser 430	acc Thr	ctg Leu	cca Pro	aat Asn	gtt Val 435	tca Ser	ttc Phe	acc Thr	atc Ile	gga Gly 440	ggg Gly	aaa Lys	agt Ser	1407
40	ttt Phe	gag Glu 445	ctg Leu	acc Thr	ctt Leu	aag Lys	gag Glu 450	tat Tyr	gtt Val	ctt Leu	cga Arg	act Thr 455	gga Gly	gaa Glu	ggc Gly	ttt Phe	1455
45	gct Ala 460	gaa Glu	gtc Val	tgc Cys	atc Ile	agt Ser 465	gga Gly	ttc Phe	atg Met	gct Ala 470	atg Met	gat Asp	gtg Val	ccg Pro	ccg Pro	cct Pro 475	1503
50	cgt Arg	ggt Gly	ccc Pro	atc Ile	tgg Trp 480	gtt Val	ctg Leu	gga Gly	gat Asp	gtg Val 485	ttc Phe	atg Met	gga Gly	gtg Val	tac Tyr 490	cac His	1551
55	acc Thr	gtg Val	ttt Phe	gat Asp 495	tat Tyr	ggt Gly	aat Asn	ctc Leu	cgg Arg 500	atg Met	ggt Gly	ttc Phe	gca Ala	aga Arg 505	gct Ala	gct Ala	1599
60	tag acaagactgt ttatttcgtc tactgtttga cggtcctaag agaagctatg																1652
65	aagacatgta gtagcttgta aattaggatt taattatgct tggctggttt atgggtgggtg																1712
	cttttaatat tatatgtaat gtaagcagat atgttacctt gtttagagt ttcaaggaaa																1772
	ctgcaatatt tacttccggt aaaaaaaaaa aaaaaaaaaa aaaaaaa																1819
	<210> 8																
	<211> 507																
	<212> PRT																
	<213> Coffea canephora																
	<400> 8																
	Met Glu Arg Arg Tyr Leu Trp Ala Ala Phe Val Leu Gly Ala Ile Val																

	1	5	10	15
5	Cys	Ser	Leu	Phe 20
	Pro	Leu	Pro	Ser
	Glu 25	Gly	Leu	Lys
	Arg	Ile 30	Ser	Leu
10	Lys	Lys	Lys 35	Pro
	Leu	Asp	Ile	Gln 40
	Ser	Ile	Arg	Ala
	Ala 45	Lys	Leu	Ala
15	His	Leu 50	Glu	Ser
	Thr	His	Gly 55	Ala
	Gly	Arg	Lys	Glu 60
	Met	Asp	Asn	Asn
20	Leu	Gly	Ser	Ser
	Asn	Glu 70	Asp	Ile
	Leu	Pro	Leu 75	Lys
	Asn	Tyr	Leu	Asp 80
25	Ala	Gln	Tyr	Tyr
	Gly 85	Glu	Ile	Gly
	Ile	Gly	Ile	Gly 90
	Thr	Pro	Pro	Gln
	Lys 95	Phe		
30	Thr	Val	Ile	Phe 100
	Asp	Thr	Gly	Ser
	Ser 105	Asn	Leu	Trp
	Val	Pro 110	Ser	Ala
35	Lys	Cys	Tyr 115	Phe
	Ser	Ile	Ala	Cys 120
	Trp	Leu	His	Ser
	Lys 125	Tyr	Lys	Ala
40	Lys	Lys 130	Ser	Ser
	Thr	Tyr	Thr 135	Ala
	Ile	Gly	Lys	Ser 140
	Cys	Ser	Ile	Arg
45	Tyr	Gly	Ser	Gly
	Ser 145	Ile 150	Ser	Gly
	Phe	Ser 155	Gln	Asp
	Asn	Val	Glu 160	
50	Val	Gly	Asp	Leu
	Val 165	Val	Lys	Asp
	Gln	Val 170	Phe	Ile
	Glu	Ala	Ser 175	Arg
55	Glu	Gly	Ser	Leu
	Thr 180	Phe	Val	Ile
	Ala 185	Lys	Phe	Asp
	Gly 190	Ile	Leu	Gly
60	Leu	Gly	Phe 195	Gln
	Glu	Ile	Ala	Val 200
	Asp	Asn	Met	Val
	Pro 205	Val	Trp	Tyr
65	Asn	Met 210	Val	Asp
	Gln	Gly	Leu 215	Val
	Asp	Glu	Gln	Val 220
	Phe	Ser	Phe	Trp
70	Leu	Asn	Arg	Asp
	Pro	Asn 230	Ala	Glu
	Asp	Gly	Gly 235	Glu
	Leu	Val	Phe	Gly 240
75	Gly	Val	Asp	Thr
	Asn 245	His	Phe	Lys
	Gly	Lys 250	His	Thr
	Tyr	Val	Pro 255	Val
80	Thr	Gln	Lys	Gly
	Tyr 260	Trp	Gln	Phe
	Lys 265	Met	Gly	Asp
	Phe	Leu 270	Ile	Gly
85	Asn	Val	Ser 275	Thr
	Gly	Phe	Cys	Glu 280
	Gly	Gly	Cys	Ala
	Ala 285	Ile	Val	Asp

5 Ser Gly Thr Ser Leu Leu Ala Gly Pro Thr Thr Val Val Thr Gln Ile  
 290 295 300  
 10 Asn His Ala Ile Gly Ala Glu Gly Val Val Ser Thr Glu Cys Lys Glu  
 305 310 315 320  
 15 Ile Val Ser Gln Tyr Gly Glu Leu Ile Trp Asp Leu Leu Val Ser Gly  
 325 330 335  
 20 Val Leu Pro Asp Arg Val Cys Lys Gln Ala Gly Leu Cys Pro Leu Arg  
 340 345 350  
 25 Gly Ala Gln His Glu Asn Ala Tyr Ile Lys Ser Val Val Asp Glu Glu  
 355 360 365  
 30 Asn Lys Glu Glu Ala Ser Val Gly Glu Ser Pro Met Cys Thr Ala Cys  
 370 375 380  
 35 Glu Met Ala Val Val Trp Met Gln Asn Gln Leu Lys Gln Gln Gly Thr  
 385 390 395 400  
 40 Lys Glu Lys Val Leu Ala Tyr Val Asn Gln Leu Cys Glu Ser Ile Pro  
 405 410 415  
 45 Ser Pro Met Gly Glu Ser Ile Ile Asp Cys Asn Ser Leu Ser Thr Leu  
 420 425 430  
 50 Pro Asn Val Ser Phe Thr Ile Gly Gly Lys Ser Phe Glu Leu Thr Leu  
 435 440 445  
 55 Lys Glu Tyr Val Leu Arg Thr Gly Glu Gly Phe Ala Glu Val Cys Ile  
 450 455 460  
 60 Ser Gly Phe Met Ala Met Asp Val Pro Pro Pro Arg Gly Pro Ile Trp  
 465 470 475 480  
 Val Leu Gly Asp Val Phe Met Gly Val Tyr His Thr Val Phe Asp Tyr  
 485 490 495  
 Gly Asn Leu Arg Met Gly Phe Ala Arg Ala Ala  
 500 505



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**